Stochastic Loss Reserving Using Bayesian MCMC Models

Glenn Meyers – FCAS, MAAA, CERA, Ph.D.

ASTIN Colloquium

May 23, 2013

Background

- Risk based capital proposals, e.g. EU Solvency II and USA SMI rely on stochastic models.
  - VaR@99.5% and TVaR@99%
- There are many stochastic loss reserve models that claim to predict the distribution of ultimate losses.

How good are these models?

- This presentation describes tests of the predictions of currently popular stochastic loss reserve models on real data from 50 insurers in each of four lines of insurances.
- It proposes two new models that improve the predictions.

The CAS Loss Reserve Database Created by Meyers and Shi

With Permission of American NAIC

- Schedule P (Data from Parts 1-4) for several US Insurers
  - Private Passenger Auto
  - Commercial Auto
  - Workers' Compensation
  - General Liability
  - Product Liability
  - Medical Malpractice (Claims Made)
- Available on CAS Website

  http://www.casact.org/research/index.cfm?fa=loss_reserves_data

Notation

- w = Accident Year $w = 1,\ldots,10$
- d = Development Year $d = 1,\ldots,10$
- $C_{w,d}$ = Cumulative (either incurred or paid) loss
- $I_{w,d}$ = Incremental paid loss $= C_{w,d} - C_{w-1,d}$

Illustrative Insurer – Incurred Losses

		Cumulative Incurred Losses										
Premium	AY/Lag	1	2	3	4	5	6	7	8	9	10	Source
5812	1988	1722	3830	3603	3835	3873	3895	3918	3918	3917	3917	1997
4908	1989	1581	2192	2528	2533	2528	2530	2534	2541	2538	2532	1998
5454	1990	1834	3009	3488	4000	4105	4087	4112	4170	4271	4279	1999
5165	1991	2305	3473	3713	4018	4295	4334	4343	4340	4342	4341	2000
5214	1992	1832	2625	3086	3493	3521	3563	3542	3541	3541	3587	2001
5230	1993	2289	3160	3154	3204	3190	3206	3351	3289	3267	3268	2002
4992	1994	2881	4254	4841	5176	5551	5689	5683	5688	5684	5684	2003
5466	1995	2489	2956	3382	3755	4148	4123	4126	4127	4128	4128	2004
5226	1996	2541	3307	3789	3973	4031	4157	4143	4142	4144	4144	2005
4962	1997	2203	2934	3608	3977	4040	4121	4147	4155	4183	4181	2006

Illustrative Insurer – Paid Losses

		Cumulative Paid Losses										
Premium	AY/Lag	1	2	3	4	5	6	7	8	9	10	Source
5812	1988	952	1529	2813	3647	3724	3832	3899	3907	3911	3912	1997
4908	1989	849	1564	2202	2432	2468	2487	2513	2526	2531	2527	1998
5454	1990	983	2211	2830	3832	4039	4065	4102	4155	4268	4274	1999
5165	1991	1657	2685	3169	3600	3900	4320	4332	4338	4341	4341	2000
5214	1992	932	1940	2626	3332	3368	3491	3531	3540	3540	3583	2001
5230	1993	1162	2402	2799	2996	3034	3042	3230	3238	3241	3268	2002
4992	1994	1478	2980	3945	4714	5462	5680	5682	5683	5684	5684	2003
5466	1995	1240	2080	2607	3080	3678	4116	4117	4125	4128	4128	2004
5226	1996	1326	2412	3367	3843	3965	4127	4133	4141	4142	4144	2005
4962	1997	1413	2683	3173	3674	3805	4005	4020	4095	4132	4139	2006

Criteria for a "Good" Stochastic Loss Reserve Model

- Using the upper triangle "training" data, predict the distribution of the outcomes in the lower triangle
  - Can be observations from individual (AY, Lag) cells or sums of observations in different (AY,Lag) cells.
- Using the predictive distributions, find the percentiles of the outcome data.
- The percentiles should be uniformly distributed.
  - Histograms
  - Test with PP Plots/KS tests
    - Plot Expected vs Predicted Percentiles
    - KS 95% critical values = 19.2 for n = 50 and 9.6 for n = 200

Illustrative Tests of Uniformity



Data Used in Study

- Insurers listed in Meyers – Summer 2012 e-Forum
- 50 Insurers from four lines of business
  - Commercial Auto
  - Personal Auto
  - Workers' Compensation
  - Other Liability
- Both paid and incurred losses

Test of Mack Model on Incurred Data



Conclusion – The Mack model predicts tails that are too light.

Test of Mack Model on Paid Data



Conclusion – The Mack model is biased upward.

Test of Bootstrap ODP on Paid Data



Conclusion – The Bootstrap ODP model is biased upward.

Possible Responses to the model failures

- The “Black Swans” got us again!
  - We do the best we can in building our models, but the real world keeps throwing curve balls at us.
  - Every few years, the world gives us a unique “black swan” event.
- Build a better model.
  - Use a model, or data, that sees the “black swans.”

Bayesian MCMC Models

- Use R and JAGS (Just Another Gibbs Sampler) packages
- Get a sample of 10,000 parameter sets from the posterior distribution of the model
- Use the parameter sets to get 10,000 simulated outcomes
- Calculate summary statistics of the simulated outcomes
  - Mean
  - Standard deviation
  - Percentile of the actual outcome

The Correlated Chain Ladder (CCL) Model

- $\mu_{1,d} = \alpha_1 + \beta_d$
- $C_{1,d} \sim \text{lognormal}(\boldsymbol{\mu_{1,d}}, \sigma_d)$
- $\mu_{w,d} = \alpha_w + \beta_d + \rho\cdot(\log(C_{w-1,d}) - \boldsymbol{\mu_{w-1,d}})$ for $w = 2,\ldots,10$
- $C_{w,d} \sim \text{lognormal}(\boldsymbol{\mu_{w,d}}, \sigma_d)$
- $\rho \sim U(-1,1)$
- α_w and β_d are widely distributed. $\beta_1 = 0$.
- $\sigma_d = \sum_{i=d}^{10} a_i \quad a_i \sim U(0,1)$ Forces σ_d to decrease as d increases
- Estimate distribution of $\sum_{w=1}^{10} C_{w,10}$

The Correlated Chain Ladder Model Predicts Distributions with Thicker Tails

- Chain ladder applies factors to last ***fixed*** observation
- CCL uses ***uncertain*** "level" parameters for each accident year.
  - $Var\left[C_{w,d}\right] = E_{\alpha_w}\left[Var\left[C_{w,d} \mid \alpha_w\right]\right] + Var_{\alpha_w}\left[E\left[C_{w,d} \mid \alpha_w\right]\right]$
- Mack uses point estimations of parameters
- CCL uses Bayesian estimation to get a posterior distribution of parameters
- Mack assumes independence between accident years
- CCL allows for correlation between accident years
  - $Corr[\log(C_{w\text{-}1,d}),\log(C_{w,d})] = \rho$

Posterior Distribution of ρ for Illustrative Insurer



Generally Positive Posterior Means of ρ



Predicting the Distribution of Outcomes

- Use JAGS (**J**ust **A**nother **G**ibbs **S**ampler) software to produce a sample of 10,000 $\{\alpha_w\}$, $\{\beta_d\}$,$\{\sigma_d\}$ and $\{\rho\}$ from the posterior distribution.
- For each member of the sample
  - $\mu_1 = \alpha_1 + \beta_{10}$
  - For w = 2 to 10
    - $C_{w,10}$ = random lognormal $(\alpha_w + \beta_{10} + \rho\cdot(\log(C_{w\text{-}1,10}) - \mu_{w\text{-}1})),\sigma_d)$
  - Calculate $\sum_{w=1}^{10} C_{w,10}$
- Calculate summary statistics, e.g. $E\left[\sum_{w=1}^{10} C_{w,10}\right]$ and $Var\left[\sum_{w=1}^{10} C_{w,10}\right]$

Results for the Illustrative Incurred Data

	CCL				Mack				Outcome
w	C_{w,10}	SD	CV		C_{w,10}	SD	CV		C_{w,10}
1	3,917	0	0.0000		3,917	0	0.0000		3,917
2	2,546	62	0.0244		2,538	0	0.0000		2,532
3	4,111	119	0.0289		4,167	3	0.0007		4,279
4	4,316	136	0.0315		4,367	37	0.0085		4,341
5	3,552	126	0.0355		3,597	34	0.0095		3,587
6	3,321	150	0.0452		3,236	40	0.0124		3,268
7	5,285	295	0.0558		5,358	146	0.0272		5,684
8	3,805	335	0.0880		3,765	225	0.0598		4,128
9	4,180	615	0.1471		4,013	412	0.1027		4,144
10	4,141	1,371	0.3311		3,955	878	0.2220		4,181
Total	39,174	1,869	0.0477		38,914	1,057	0.0272		40,061
Percentile		73.40				86.03			

Note the increase in the standard error of CCL over Mack.

Compare SDs for All 200 Triangles



Test of Mack Model on Incurred Data



Conclusion – The Mack model predicts tails that are too light.

Test of CCL on Incurred Data



Conclusion – CCL model percentiles lie within KS statistical bounds.

Improvement with Incurred Data

- Accomplished by “pumping up” the variance of Mack model.

What About Paid Data?

- Start by looking at CCL model on cumulative paid data.

Test of Bootstrap ODP on Paid Data



Conclusion – The Bootstrap ODP model is biased upward.

Test of CCL on Paid Data



Conclusion – Roughly the same performance a bootstrapping and Mack

How Do We Correct the Bias?

- Look at models with payment year trend.
  - Ben Zehnwirth has been championing these for years.
- Payment year trend does not make sense with cumulative data!
  - Settled claims are unaffected by trend.
- Recurring problem with incremental data – Negatives!
  - We need a skewed distribution that has support over the entire real line.

The Lognormal-Normal (ln-n) Mixture

$X \sim \text{Normal}(Z, \delta)$, $Z \sim \text{Lognormal}(\mu, \sigma)$



The Correlated Incremental Trend (CIT) Model

- $\mu_{w,d} = \alpha_w + \beta_d + \tau\cdot(w + d - 1)$
- $Z_{w,d} \sim \text{lognormal}(\mu_{w,d}, \sigma_d)$ subject to $\sigma_1 < \sigma_2 < \ldots < \sigma_{10}$
- $I_{1,d} \sim \text{normal}(Z_{1,d}, \delta)$
- $I_{w,d} \sim \text{normal}(Z_{w,d} + \rho\cdot(I_{w-1,d} - Z_{w-1,d})\cdot e^{\tau}, \delta)$
- Estimate the distribution of $\sum_{w=1}^{10} C_{w,10}$
- "Sensible" priors on α_w, σ_d, and τ. $\beta_1 = 0$
  - Needed to control σ_d
  - Interaction between τ, α_w and β_d.

CIT Model for Illustrative Insurer

	CIT				CCL				Outcome
w	$C_{w,10}$	SD	CV		$C_{w,10}$	SD	CV		$C_{w,10}$
1	3912	0	0		3912	0	0.0000		3912
2	2536	5	0.002		2563	110	0.0429		2527
3	4175	11	0.0026		4153	189	0.0455		4274
4	4378	29	0.0066		4320	224	0.0519		4341
5	3539	35	0.0099		3570	207	0.0580		3583
6	3043	105	0.0345		3403	255	0.0749		3268
7	5037	114	0.0226		5207	465	0.0893		5684
8	3501	556	0.1588		3649	467	0.1280		4128
9	3980	710	0.1784		4409	895	0.2030		4144
10	4661	1484	0.3184		5014	2435	0.4856		4139
Total	38763	1803	0.0465		40200	3070	0.0764		40000
Percentile		81.87				51.24			

Posterior Distribution of μ and τ for Illustrative Insurer





Posterior Mean ρ for All Insurers On Paid Data



Posterior Mean ρ for All Insurers On Incurred Data



Posterior Mean τ for All Insurers



Test of Bootstrap ODP on Paid Data



Conclusion – The Bootstrap ODP model is biased upward.

Test of CIT with ρ = 0 on Paid Data



Conclusion – Overall improvement but look at Personal Auto

Test of CIT on Paid Data



Conclusion – CIT model percentiles are an improvement but do not lie within the KS bounds.

Summary

- Mack underpredicts the variability of outcomes with incurred data.
- Both Mack and Bootstrap ODP are biased high with paid data.
- Bayesian MCMC models
  - Easily modified to produce new models.
  - Easily implemented to produce predictive distributions of outcomes.
- CCL model improves significantly on predictions with incurred data.
  - Important feature – Correlation between accident years
- CIT models improves somewhat on predictions with paid data.
  - Important features – Payment year trend and correlation between accident years
- Shortcoming – Study needs to be repeated on different time periods.

Purpose of CAS Loss Reserve Database

- Large scale testing of model predictions.